

Patient Information	Specimen Information	Client Information
TEST, PATIENT DOB: 01/01/1969 AGE: 45 Gender: F Fasting: Y Phone: NG Patient ID: NG	Specimen: CW014217A Requisition: 0046280 Collected: 04/28/2014 / 10:00 EDT Received: 04/28/2014 / 12:54 EDT Reported: 04/28/2014 / 17:59 EDT (* A Copy Sent To)	Client #: 97502840 00000000 DOE, JANE TEST CLIENT (HQ) NEL Attn: TEST DEPARTMENT 400 EGYPT RD # 2009 1/3 NORRISTOWN, PA 19403-3406

Diabetes Risk Panel with Score


Glucose value and assessment

Lab: QWA




HbA1c value and assessment

Lab: QWA



8-Year diabetes risk (%)

Lab: QWA



The APPROXIMATE 8-year Percentage Risk estimate is based upon the assessment of adults ages 30-79 in the Framingham study (Wilson PW, Meigs JB, Sullivan L, Fox CS, Nathan DM, D'Agostino RB Sr. 2007 Prediction of incident diabetes mellitus in middle-aged adults: the Framingham Offspring Study Arch Intern Med. 167:1068-74). An estimate will not be provided if HbA1c and/or glucose indicate that the patient has diabetes. If fasting conditions were not met, use caution when interpreting the glucose test result and the 8-year Percentage Risk.

Patient Information	Specimen Information	Client Information
TEST, PATIENT DOB: 01/01/1969 AGE: 45 Gender: F Fasting: Y Patient ID: NG	Specimen: CW014217A Collected: 04/28/2014 / 10:00 EDT Received: 04/28/2014 / 12:54 EDT Reported: 04/28/2014 / 17:59 EDT (* A Copy Sent To)	Client #: 97502840 DOE, JANE

Test values
Lab: QWA

Test Name	Result	Reference Range/Comments
Glucose	115 H	65-99 mg/dL See Note 1
Hemoglobin A1c	6.1 H	<5.7 % of total Hgb See Note 2
Total Cholesterol	164	125-200 mg/dL
HDL Cholesterol	81	> OR = 46 mg/dL
Triglycerides	45	<150 mg/dL
LDL Cholesterol (Calculated)	74	<130 mg/dL (calc) See Note 3
Non-HDL Cholesterol	83	mg/dL (calc) See Note 4
Cholesterol/HDL Ratio (Calculated)	2.0	< OR = 5.0 (calc)

Personal factors
Lab: QWA

Component	Result	Component	Result
HEIGHT FEET	5 ft	SYSTOLIC BLOOD PRESSURE	106 mmHg
HEIGHT INCHES	4 in	DIASTOLIC BLOOD PRESSURE	65 mmHg
WEIGHT	150 lbs	TREATMENT FOR HIGH B.P.	NO
CALCULATED BMI	25.7	PARENTAL HISTORY OF DIAB	YES

Note 1

Fasting reference interval

Note 2

According to ADA guidelines, hemoglobin A1c <7.0% represents optimal control in non-pregnant diabetic patients. Different metrics may apply to specific patient populations. Standards of Medical Care in Diabetes-2013. Diabetes Care. 2013;36:s11-s66

For the purpose of screening for the presence of diabetes

<5.7% Consistent with the absence of diabetes

5.7-6.4% Consistent with increased risk for diabetes (prediabetes)

>or=6.5% Consistent with diabetes

This assay result is consistent with a higher risk of diabetes.

Currently, no consensus exists for use of hemoglobin A1c for diagnosis of diabetes for children.

Note 3

Desirable range <100 mg/dL for patients with CHD or

Patient Information	Specimen Information	Client Information
TEST, PATIENT DOB: 01/01/1969 AGE: 45 Gender: F Fasting: Y Patient ID: NG	Specimen: CW014217A Collected: 04/28/2014 / 10:00 EDT Received: 04/28/2014 / 12:54 EDT Reported: 04/28/2014 / 17:59 EDT (* A Copy Sent To)	Client #: 97502840 DOE, JANE

diabetes and <70 mg/dL for diabetic patients with known heart disease.

Note 4

Target for non-HDL cholesterol is 30 mg/dL higher than LDL cholesterol target.

PERFORMING SITE:

QWA QUEST DIAGNOSTICS WALLINGFORD-CL 0091, 3 STERLING DRIVE, WALLINGFORD, CT 06492 Laboratory Director: NEENA SINGH, MD, CLIA: 07D0093126

* Copy To Client: HQTEST - FAX AS FINAL